




PubMed



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed	for		Go		Clear			
About Entrez	Limits	Preview/Index	History	Clipboard	Details			
Text Version	Display	Abstract	Show: 20	Sort	Send to	Text		

1: Plast Reconstr Surg. 2002 Jun;109(7):2384-97.

Related Articles, Links



Hypoxia and VEGF up-regulate BMP-2 mRNA and protein expression in microvascular endothelial cells: implications for fracture healing.

Bouletreau PJ, Warren SM, Spector JA, Peled ZM, Gerrets RP, Greenwald JA, Longaker MT.

Department of Surgery, Stanford University School of Medicine, 257 Campus Drive, Stanford, CA 94305-5148, USA.

The endothelium is a metabolically active secretory tissue, capable of responding to a wide array of environmental stimuli. Hypoxia and vascular endothelial growth factor (VEGF) are two components of the putative fracture microenvironment. This study investigated the role of hypoxia and VEGF on endothelial cell activation as it relates to the bone repair process. It was hypothesized that endothelial cells may have an important osteogenic role in fracture healing through the production of bone morphogenetic protein-2 (BMP-2), an osteogenic cytokine at the fracture site. Therefore, BMP-2 mRNA and protein expression in endothelial cells under hypoxia and/or VEGF treatment was studied. The authors observed a 2-fold to 3-fold up-regulation of BMP-2 mRNA expression in bovine capillary endothelial cells and human microvascular endothelial cells stimulated with hypoxia or rhVEGF. Furthermore, the combined effects of hypoxia and rhVEGF appeared to be additive on BMP-2 mRNA expression in bovine capillary endothelial cells. Actinomycin D and cycloheximide studies suggested that the increased mRNA expression was transcriptionally regulated. BMP-2 protein expression was up-regulated after 24 and 48 hours of treatment with either hypoxia or rhVEGF in bovine capillary endothelial cells. Surprisingly, the data suggest that endothelial cells may play not only an angiogenic role but also an osteogenic role by a direct stimulation of the osteoblasts, through the enhanced expression of a potent osteogenic factor, BMP-2, at the fracture site.

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central
Privacy Policy

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MESH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

PMID: 12045566 [PubMed - indexed for MEDLINE]

Display	Abstract	Show: 20	Sort	Send to	Text
---------	----------	----------	------	---------	------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Aug 6 2003 12:56:11

Best Available Copy